

নিচের যোগজগুলির মান নির্ণয় কর :

1.(a) $\int \frac{1}{x} \left(x + \frac{1}{x}\right) dx$ [স. '০৫]

$$= \int (1 + x^{-2}) dx = x + \frac{x^{-2+1}}{-2+1} + c$$

$$= x - \frac{1}{x} + c$$

1(b) $\int \frac{(e^x + 1)^2}{\sqrt{e^x}} dx$ [স. '০২]

$$= \int \frac{e^{2x} + 2e^x + 1}{e^{\frac{x}{2}}} dx$$

$$= \int (e^{2x - \frac{x}{2}} + 2e^{x - \frac{x}{2}} + e^{-\frac{x}{2}}) dx$$

$$= \int (e^{\frac{3x}{2}} + 2e^{\frac{x}{2}} + e^{-\frac{x}{2}}) dx$$

$$= \frac{e^{\frac{3x}{2}}}{\frac{3}{2}} + 2 \frac{e^{\frac{x}{2}}}{\frac{1}{2}} + \frac{e^{-\frac{x}{2}}}{-\frac{1}{2}} + c$$

$$= \frac{2}{3} e^{\frac{3x}{2}} + 4e^{\frac{x}{2}} - 2e^{-\frac{x}{2}} + c$$

1(c) $\int (1 + x^{-1} + x^{-2}) dx$ [স. '০৯]

$$= \int \left(1 + \frac{1}{x} + x^{-2}\right) dx$$

$$= x + \ln x + \frac{x^{-2+1}}{-2+1} + c = x + \ln x - x^{-1} + c$$

নিয়ম : হরের অনুবন্ধি রাশি দ্বারা লব ও হরকে গুণ করে হরকে $\sqrt{\quad}$ যুক্ত করতে হয়।

2.(a) $\int \frac{1}{\sqrt{x} - \sqrt{x-1}} dx$

$$= \int \frac{\sqrt{x} + \sqrt{x-1}}{(\sqrt{x} - \sqrt{x-1})(\sqrt{x} + \sqrt{x-1})} dx$$

$$= \int \frac{\sqrt{x} + \sqrt{x-1}}{x - (x-1)} dx = \int \frac{\sqrt{x} + \sqrt{x-1}}{x - x + 1} dx$$

$$= \int \{x^{\frac{1}{2}} + (x-1)^{\frac{1}{2}}\} dx$$

$$= \frac{x^{\frac{1}{2}+1}}{\frac{1}{2}+1} + \frac{(x-1)^{\frac{1}{2}+1}}{\frac{1}{2}+1} + c$$

$$= \frac{2}{3} [x^{3/2} + (x-1)^{3/2}] + c$$

2(b) $\int \frac{dx}{\sqrt{x+1} + \sqrt{x-1}}$ [স. '০২; দি. '১০]

$$= \int \frac{\sqrt{x+1} - \sqrt{x-1}}{(\sqrt{x+1} + \sqrt{x-1})(\sqrt{x+1} - \sqrt{x-1})} dx$$

$$= \int \frac{\sqrt{x+1} - \sqrt{x-1}}{(x+1) - (x-1)} dx$$

$$= \frac{1}{2} \int [(x+1)^{1/2} - (x-1)^{1/2}] dx$$

$$= \frac{1}{2} \left[\frac{(x+1)^{\frac{1}{2}+1}}{\frac{1}{2}+1} - \frac{(x-1)^{\frac{1}{2}+1}}{\frac{1}{2}+1} \right] + c$$

$$= \frac{1}{2} \left[\frac{(x+1)^{\frac{3}{2}}}{\frac{3}{2}} - \frac{(x-1)^{\frac{3}{2}}}{\frac{3}{2}} \right] + c$$

$$= \frac{1}{3} [(x+1)^{\frac{3}{2}} - (x-1)^{\frac{3}{2}}] + c \text{ (Ans.)}$$

3.(a) $\int \frac{dx}{1 - \sin x}$ [স. '০৭]

$$= \int \frac{(1 + \sin x) dx}{(1 - \sin x)(1 + \sin x)}$$

$$= \int \frac{(1 + \sin x) dx}{1 - \sin^2 x} = \int \frac{(1 + \sin x) dx}{\cos^2 x}$$

$$= \int \left(\frac{1}{\cos^2 x} + \frac{\sin x}{\cos^2 x} \right) dx$$

$$= \int (\sec^2 x + \sec x \tan x) dx$$

$$= \tan x + \sec x + c$$

3(b) $\int \frac{dx}{1 + \sin x}$ [য.'০৭, '১৩; চ.'১০ প্র.ভ.প.'০৩]

$$= \int \frac{(1 - \sin x) dx}{(1 + \sin x)(1 - \sin x)}$$

$$= \int \frac{(1 - \sin x) dx}{1 - \sin^2 x} = \int \frac{(1 - \sin x) dx}{\cos^2 x}$$

$$= \int \left(\frac{1}{\cos^2 x} - \frac{\sin x}{\cos^2 x} \right) dx$$

$$= \int (\sec^2 x - \sec x \tan x) dx$$

$$= \tan x - \sec x + c$$

3(c) $\int \frac{dx}{1 + \cos 2x}$ [কু.'০৮]

$$= \int \frac{dx}{2 \cos^2 x} = \frac{1}{2} \int \sec^2 x dx = \frac{1}{2} \tan x + c$$

3(d) $\int \sqrt{1 + \cos x} dx$ [প্র.ভ.প.'০৪]

$$= \int \sqrt{2 \cos^2 \frac{x}{2}} dx = \int \sqrt{2} \cos \frac{x}{2} dx$$

$$= 2\sqrt{2} \int \cos \frac{x}{2} d\left(\frac{x}{2}\right) \quad \left[\because d\left(\frac{x}{2}\right) = \frac{1}{2} dx \right]$$

$$= 2\sqrt{2} \sin \frac{x}{2} + c$$

3(e) $\int \sqrt{1 - \cos 2x} dx$ [চ.'০৫, '০৯; সি.'০৬; ব.'০৮]

$$= \int \sqrt{2 \sin^2 x} dx = \int \sqrt{2} \sin x dx$$

$$= \sqrt{2}(-\cos x) + c = -\sqrt{2} \cos x + c$$

3(f) $\int \sqrt{1 - \cos 4x} dx$ [চ.'০৭]

$$= \int \sqrt{2 \sin^2 2x} dx = \int \sqrt{2} \sin 2x dx$$

$$= \sqrt{2} \left(-\frac{\cos 2x}{2} \right) + c = -\frac{1}{\sqrt{2}} \cos 2x + c$$

3(g) $\int \sec x (\sec x - \tan x) dx$ [ব.'১৩]

$$= \int (\sec^2 x - \sec x \tan x) dx$$

$$= \tan x - \sec x + c$$

4(a) $\int \sqrt{1 - \sin 2x} dx$

$$= \int \sqrt{\sin^2 x + \cos^2 x - 2 \sin x \cos x} dx$$

$$= \int \sqrt{(\sin x - \cos x)^2} dx$$

$$= \int (\sin x - \cos x) dx \text{ বা } \int (\cos x - \sin x) dx$$

$$= -\cos x - \sin x + c \text{ বা } \sin x + \cos x + c$$

4(b) $\int \frac{\cos 2x}{\sqrt{1 - \sin 2x}} dx$

$$= \int \frac{\cos^2 x - \sin^2 x}{\sqrt{\sin^2 x + \cos^2 x - 2 \sin x \cos x}} dx$$

$$= \int \frac{\cos^2 x - \sin^2 x}{\sqrt{(\sin x - \cos x)^2}} dx$$

$$= \int \frac{(\cos x - \sin x)(\cos x + \sin x)}{\cos x - \sin x} dx$$

$$\text{বা, } \int \frac{(\cos x - \sin x)(\cos x + \sin x)}{\sin x - \cos x} dx$$

$$= \int (\cos x + \sin x) dx \text{ বা, } - \int (\cos x + \sin x) dx$$

$$= \sin x - \cos x \text{ বা, } -(\sin x - \cos x)$$

4(c) $\int (\sin x + \cos x)^2 dx$ [প্র.ভ.প.'১০]

$$= \int (\sin^2 x + \cos^2 x + 2 \sin x \cos x) dx$$

$$= \int (1 + \sin 2x) dx = x - \frac{1}{2} \cos 2x + c$$

5(a) $\int \sin 5x \sin 3x dx$ [ব.'০৮, '১২; য.'১০; চ.'১২]

$$= \int \frac{1}{2} \{ \cos(5x - 3x) - \cos(5x + 3x) \} dx$$

$$= \frac{1}{2} \int (\cos 2x - \cos 8x) dx$$

$$= \frac{1}{2} \left(\frac{\sin 2x}{2} - \frac{\sin 8x}{8} \right) + c$$

$$= \frac{1}{4} \sin 2x - \frac{1}{16} \sin 8x + c$$

$$5(b) \int \sin 4x \sin 2x dx \quad [\text{য. '০৪; রা. '০৫; দি. '১১}]$$

$$= \int \frac{1}{2} \{ \cos(4x - 2x) - \cos(4x + 2x) \} dx$$

$$= \frac{1}{2} \int (\cos 2x - \cos 6x) dx$$

$$= \frac{1}{2} \left(\frac{\sin 2x}{2} - \frac{\sin 6x}{6} \right) + c$$

$$= \frac{1}{4} \sin 2x - \frac{1}{12} \sin 6x + c$$

$$5(c) \int 3 \sin 3x \cos 4x dx \quad [\text{সি. '০২, '০৩; ব. '০৬, '১০}]$$

$$= \int \frac{3}{2} \{ \sin(4x + 3x) - \sin(4x - 3x) \} dx$$

$$= \frac{3}{2} \int (\sin 7x - \sin x) dx$$

$$= \frac{3}{2} \left(-\frac{1}{7} \cos 7x + \cos x \right) + c$$

$$= \frac{3}{14} (7 \cos x - \cos 7x) + c$$

$$5(d) \int \sin 3x \cos 5x dx \quad [\text{ক. '০৬; সি., দি. '১২}]$$

$$= \int \frac{1}{2} \{ \sin(5x + 3x) - \sin(5x - 3x) \} dx$$

$$= \int \frac{1}{2} (\sin 8x - \sin 2x) dx$$

$$= \frac{1}{2} \left(-\frac{1}{8} \cos 8x + \frac{1}{2} \cos 2x \right) + c$$

$$= \frac{1}{16} (4 \cos 2x - \cos 8x) + c$$

$$5(e) \int 4 \cos 4x \sin 5x dx \quad [\text{রা. '০৩}]$$

$$= \int 2 \{ \sin(5x + 4x) + \sin(5x - 4x) \} dx$$

$$= \int 2 (\sin 9x + \sin x) dx$$

$$= 2 \left(-\frac{1}{9} \cos 9x - \cos x \right) + c$$

$$= -\frac{2}{9} (\cos 9x + 9 \cos x) + c$$

$$5(f) \int 5 \cos 5x \sin 4x dx \quad [\text{ঢা. '০৬; দি., সি. '১৪}]$$

$$= \int \frac{5}{2} \{ \sin(5x + 4x) - \sin(5x - 4x) \} dx$$

$$= \int \frac{5}{2} (\sin 9x - \sin x) dx$$

$$= \frac{5}{2} \left(-\frac{1}{9} \cos 9x + \cos x \right) + c$$

$$= \frac{5}{18} (9 \cos x - \cos 9x) + c$$

$$5(g) \int \sin px \cos qx dx, (p > q)$$

[ঢা. '০৩; সি. '০৭]

$$= \int \frac{1}{2} \{ \sin(p+q)x + \sin(p-q)x \} dx$$

$$= \frac{1}{2} \left\{ -\frac{\cos(p+q)x}{p+q} - \frac{\cos(p-q)x}{p-q} \right\} + c$$

$$= -\frac{1}{2} \left\{ \frac{\cos(p+q)x}{p+q} + \frac{\cos(p-q)x}{p-q} \right\} + c$$

$$6(a) \int \cos^2 x dx \quad [\text{ঢ. '০৮}]$$

$$= \int \frac{1}{2} (1 + \cos 2x) dx = \frac{1}{2} \left\{ \int dx + \int \cos 2x dx \right\}$$

$$= \frac{1}{2} \left(x + \frac{\sin 2x}{2} \right) + c$$

$$6(b) \int \cos^2 2x dx \quad [\text{ঢা. '০০}]$$

$$= \int \frac{1}{2} (1 + \cos 4x) dx = \frac{1}{2} \left(x + \frac{\sin 4x}{4} \right) + c$$

$$6(c) \int (2 \cos x + \sin x) \cos x dx \quad [\text{ঢা. '০৫}]$$

$$= \int (2 \cos^2 x + \sin x \cos x) dx$$

$$= \int \left(1 + \cos 2x + \frac{1}{2} \sin 2x \right) dx$$

$$= x + \frac{1}{2} \sin 2x + \frac{1}{2} \left(-\frac{1}{2} \cos 2x \right) + c$$

$$= x + \frac{1}{2} \sin 2x - \frac{1}{4} \cos 2x + c$$

6(d) $\int \sin^3 2x dx$

$= \int \frac{1}{4} (3 \sin 2x - \sin 6x) dx$

$= \frac{1}{4} \left\{ 3 \cdot \left(-\frac{1}{2} \cos 2x\right) + \frac{1}{6} \cos 6x \right\} + c$

$= \frac{1}{8} (-3 \cos 2x + \frac{1}{3} \cos 6x) + c$

6(e) $\int \sin^4 x dx$

$\sin^4 x dx = (\sin^2 x)^2 = \left\{ \frac{1}{2} (1 - \cos 2x) \right\}^2$

$= \frac{1}{4} \{ 1 - 2 \cos 2x + \cos^2 2x \}$

$= \frac{1}{4} \left\{ 1 - 2 \cos 2x + \frac{1}{2} (1 + \cos 4x) \right\}$

$= \frac{1}{4} \left[1 - 2 \cos 2x + \frac{1}{2} + \frac{1}{2} \cos 4x \right]$

$= \frac{1}{4} \left[\frac{3}{2} - 2 \cos 2x + \frac{1}{2} \cos 4x \right]$

$\therefore \int \sin^4 x dx$

$= \frac{1}{4} \left(\frac{3}{2} x - 2 \cdot \frac{1}{2} \sin 2x + \frac{1}{2} \cdot \frac{1}{4} \sin 4x \right) + c$

$= \frac{1}{4} \left(\frac{3}{2} x - \sin 2x + \frac{1}{8} \sin 4x \right) + c$

6(f) $\int \cos^4 x dx$ [রা.'০৭, '১৪; সি.'০৮; দি.'১৩; গ.'১৪]

$\cos^4 x dx = (\cos^2 x)^2 = \left\{ \frac{1}{2} (1 + \cos 2x) \right\}^2$

$= \frac{1}{4} \{ 1 + 2 \cos 2x + \cos^2 2x \}$

$= \frac{1}{4} \left\{ 1 + 2 \cos 2x + \frac{1}{2} (1 + \cos 4x) \right\}$

$= \frac{1}{4} \left[1 + 2 \cos 2x + \frac{1}{2} + \frac{1}{2} \cos 4x \right]$

$= \frac{1}{4} \left[\frac{3}{2} + 2 \cos 2x + \frac{1}{2} \cos 4x \right]$

$\therefore \int \cos^4 x dx$

[ঢা.'০১]

$= \int \frac{1}{4} \left(\frac{3}{2} + 2 \cos 2x + \frac{1}{2} \cos 4x \right) dx$

$= \frac{1}{4} \left(\frac{3}{2} x + 2 \cdot \frac{1}{2} \sin 2x + \frac{1}{2} \cdot \frac{1}{4} \sin 4x \right) + c$

$= \frac{1}{4} \left(\frac{3}{2} x + \sin 2x + \frac{1}{8} \sin 4x \right) + c$ (Ans.)

সম্ভাব্য ধাপসহ প্রশ্ন:

নিচের যোগজগুলি মান নির্ণয় কর :

7(a) $\int \frac{4(\sqrt[3]{x^2} + 4)^2}{3\sqrt[3]{x}} dx = \frac{4}{3} \int \frac{(x^{\frac{2}{3}} + 4)^2}{x^{\frac{1}{3}}} dx$

$= \frac{4}{3} \int \frac{x^{\frac{4}{3}} + 8x^{\frac{2}{3}} + 16}{x^{\frac{1}{3}}} dx$

$= \frac{4}{3} \int (x^{\frac{4}{3}-\frac{1}{3}} + 8x^{\frac{2}{3}-\frac{1}{3}} + 16x^{-\frac{1}{3}}) dx$

$= \frac{4}{3} \int (x + 8x^{\frac{1}{3}} + 16x^{-\frac{1}{3}}) dx$

$= \frac{4}{3} \left(\frac{x^2}{2} + 8 \frac{x^{\frac{1}{3}+1}}{\frac{1}{3}+1} + 16 \frac{x^{-\frac{1}{3}+1}}{-\frac{1}{3}+1} \right) + c$

$= \frac{4}{3} \left(\frac{x^2}{2} + 8 \frac{x^{\frac{4}{3}}}{\frac{4}{3}} + 16 \frac{x^{\frac{2}{3}}}{\frac{2}{3}} \right) + c$

$= \frac{2}{3} (x^2 + 12x^{4/3} + 48x^{2/3}) + c$ (২)

7(b) $\int \frac{a \cot x + b \tan^2 x - c \sin^2 x}{\sin x} dx$

$= \int \left(a \frac{\cot x}{\sin x} + b \frac{\sin^2 x}{\cos^2 x \sin x} - c \sin x \right) dx$

$= \int (a \cot x \operatorname{cosec} x + b \tan x \sec x - c \sin x) dx$

$= -a \operatorname{cosec} x + b \sec x + c \cos x + c_1$ (২)

$$8(a) \int \frac{\cos 2x - \cos 2\theta}{\cos x - \cos \theta} dx$$

$$= \int \frac{2\cos^2 x - 1 - (2\cos^2 \theta - 1)}{\cos x - \cos \theta} dx$$

$$= 2 \int \frac{\cos^2 x - \cos^2 \theta}{\cos x - \cos \theta} dx$$

$$= 2 \int \frac{(\cos x + \cos \theta)(\cos x - \cos \theta)}{\cos x - \cos \theta} dx$$

$$= 2 \int (\cos x + \cos \theta) dx$$

$$= 2(\int \cos x dx + \cos \theta \int dx)$$

$$= 2(\sin x + \cos \theta \cdot x) + c$$

$$= 2(\sin x + x \cos \theta) + c$$

$$8(b) \int (\sec x + \tan x)^2 dx$$

$$= \int (\sec^2 x + \tan^2 x + 2\sec x \tan x) dx$$

$$= \int (\sec^2 x + \sec^2 x - 1 + 2\sec x \tan x) dx$$

$$= \int (2\sec^2 x - 1 + 2\sec x \tan x) dx$$

$$= 2 \tan x - x + 2\sec x + c$$

$$9(a) \int \sqrt{1 \pm \sin x} dx$$

$$= \int \sqrt{\sin^2 \frac{x}{2} + \cos^2 \frac{x}{2} \pm 2\sin \frac{x}{2} \cos \frac{x}{2}} dx \quad (১)$$

$$= \int \sqrt{(\sin \frac{x}{2} \pm \cos \frac{x}{2})^2} dx$$

$$= \int (\sin \frac{x}{2} \pm \cos \frac{x}{2}) dx \text{ বা } \int (\cos \frac{x}{2} \pm \sin \frac{x}{2}) dx \quad (১)$$

$$= 2(-\cos \frac{x}{2} \pm \sin \frac{x}{2}) + c$$

$$\text{বা } 2(\sin \frac{x}{2} \mp \cos \frac{x}{2}) + c \quad (২)$$

$$9(b) \int \frac{\sin x + \cos x}{\sqrt{1 + \sin 2x}} dx$$

$$= \int \frac{\sin x + \cos x}{\sqrt{\sin^2 x + \cos^2 x + 2\sin x \cos x}} dx \quad (১)$$

$$= \int \frac{\sin x + \cos x}{\sqrt{(\sin x + \cos x)^2}} dx$$

$$= \int \frac{\sin x + \cos x}{\sin x + \cos x} dx = \int dx = x + c \quad (১)$$

$$(১) \quad 9(c) \int \frac{\cos x + \sin x}{\cos x - \sin x} (1 - \sin 2x) dx$$

$$= \int \frac{\cos x + \sin x}{\cos x - \sin x} (\cos x - \sin x)^2 dx \quad (১)$$

$$= \int (\cos x + \sin x)(\cos x - \sin x) dx$$

$$= \int (\cos^2 x - \sin^2 x) dx = \int \cos 2x dx \quad (১)$$

$$= \frac{1}{2} \sin 2x + c \quad (২)$$

$$9(d) \int (\sin \frac{x}{2} + \cos \frac{x}{2})^2 dx$$

$$= \int (\sin^2 \frac{x}{2} + \cos^2 \frac{x}{2} + 2\sin \frac{x}{2} \cos \frac{x}{2}) dx$$

$$= \int (1 + \sin x) dx = x - \cos x + c \quad (২)$$

$$10. \int \cos^3 x dx = \int \frac{1}{4} (3\cos x + \cos 3x) dx \quad (১)$$

$$= \frac{1}{4} (3\sin x + \frac{1}{3}\sin 3x) + c \quad (১)$$

প্রশ্নমালা X B

নিচের যোগজগুলি নির্ণয় কর :

$$1.(a) \int \frac{1}{\sqrt[3]{(1-4x)}} dx = \int \frac{1}{(1-4x)^{1/3}} dx$$

$$= \int (1-4x)^{-1/3} dx = \frac{(1-4x)^{-1/3+1}}{(-1/3+1)(-4)} + c$$

$$= \frac{(1-4x)^{2/3}}{\frac{2}{3}(-4)} + c = -\frac{3}{8}(1-4x)^{2/3} + c$$

$$1(b) \int \frac{e^{5x} + e^{3x}}{e^x + e^{-x}} dx$$

[প্র.ভ.প. '৯২]

$$= \int \frac{e^{4x}(e^x + e^{-x})}{e^x + e^{-x}} dx = \int e^{4x} dx = \frac{e^{4x}}{4} + c$$

$$1(c) \text{ ধরি, } I = \int \sin x^0 dx$$

[চ. '০৪]